


Neutrino Mixings and Unified Theories

Fu-Sin Ling
Institute for Fundamental Theory
Department of Physics, University of Florida,
Gainesville, FL, 32611, USA



Abstract

The recent neutrino data converge towards a 3ν scheme with two large and one small mixing angles. Their implications for model-building are discussed. Some possible components of a flavor symmetry over quarks and leptons are singled out. We suggest that they can be naturally embedded in a unified structure with a replicated gauge group.

1 Introduction

The last five years have proven to be incredibly rich in Neutrino Physics. Numerous experiments around the globe enabled to turn neutrino oscillations from discovery into an almost complete determination of the physical parameters. Noticeably, the confirmation of the original oscillation signals from solar and atmospheric neutrinos by terrestrial and man-made sources stands as a remarkable success^{1,2} and leads to a consistent picture of the neutrino sector.

The scheme that emerges involves the three active neutrinos, whose light mass eigenstates^a are related to the flavor eigenstates by a mixing matrix U_{MNS} (the Maki-Nakagawa-Sakata matrix)^b that contains two large and one small mixing angles. The presence of a small mixing angle θ_{13} enables to write U_{MNS} as

$$U_{MNS} \simeq \begin{pmatrix} \cos \theta_{\odot} & \sin \theta_{\odot} & s_{13}e^{-i\delta} \\ -\sin \theta_{\odot} \cos \theta_{\oplus} & \cos \theta_{\odot} \cos \theta_{\oplus} & \sin \theta_{\oplus} \\ \sin \theta_{\odot} \sin \theta_{\oplus} & -\cos \theta_{\odot} \sin \theta_{\oplus} & \cos \theta_{\oplus} \end{pmatrix} \quad (1)$$

where $\theta_{\odot} \simeq \pi/6$ and $\theta_{\oplus} \simeq \pi/4$ are the mixing angles measured by solar and atmospheric neutrinos experiments respectively, $s_{13} = \sin^2 \theta_{13}$ and δ is a possible CP violating phase. The constraint from the CHOOZ experiment³ translates into a bound for θ_{13}

$$\sin^2 \theta_{13} \lesssim 0.03 \quad (2)$$

A mild hierarchy between $\Delta m_{\odot}^2$ for solar neutrinos and $\Delta m_{\oplus}^2$ for atmospheric neutrinos is also observed^{4,5}, although the absolute scale for neutrinos masses is yet to be found. Assuming the LMA solution for solar neutrinos (combined with the KamLAND result), we have

$$\Delta m_{\odot}^2 \simeq 7 \cdot 10^{-5} \text{ eV}^2; \quad \Delta m_{\oplus}^2 \simeq 3 \cdot 10^{-3} \text{ eV}^2 \quad (3)$$

^ain the effective low-energy description with Majorana masses

^bin the charged leptons mass eigenstates basis

so that

$$\frac{\Delta m_{\odot}^2}{\Delta m_{\oplus}^2} \simeq 10^{-2} \quad (4)$$

The experimental evidences that support this picture can be summarized as follows. The SNO neutral current data⁶ proves beyond any doubt that flavor conversion of solar neutrinos is taking place. Moreover, it limits severely the amount of neutrino disappearance in sterile flavors or in extra-dimensions. The observed loss of reactor anti-neutrinos in KamLAND agrees with the solar neutrinos data, and selects the LMA region as the unique solution. Moreover, the possibility of CPT violation in the neutrino sector, although not completely ruled out, seems unlikely. Of course, the controversial claim of the LSND experiment⁷ is quite problematic in this picture, because a 4ν scheme gives only a poor fit to the data. It will need the verdict of miniBoone, meanwhile, one solution is simply to drop out the LSND result.

It appears that the Yukawa structure in the neutrino sector looks very different from that of the quark sector, where steep mass hierarchies come along with small mixing angles. From the model-building point of view, the latter situation is natural, because a hierarchical structure in a mass matrix will precisely yield hierarchical masses and small mixing angles. For example, with two flavors, a mass matrix with the structure (the $\mathcal{O}(1)$ prefactor coefficients are dropped)

$$\begin{pmatrix} \epsilon^2 & \epsilon \\ \epsilon & 1 \end{pmatrix} \quad (5)$$

with $\epsilon \ll 1$ gives two eigenmasses $m_1 \sim \mathcal{O}(1)$ and $m_2 \sim \mathcal{O}(\epsilon^2)$, and a mixing angle $\theta \sim \mathcal{O}(\epsilon)$. On the contrary, hierarchical masses with large mixings lead to a mass matrix without hierarchical structure, but with correlated entries. Again, with two flavors, if $m_1 \sim \mathcal{O}(1)$, $m_2 \sim \mathcal{O}(\epsilon^2)$ and $\theta = \pi/4$ we get a matrix of the form

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \quad (6)$$

whose determinant is $D \sim \mathcal{O}(\epsilon^2)$. Of course, the absolute mass scale of neutrinos is not known yet, but naturalness arguments in unified theories favor hierarchical masses over almost degenerate masses with hyperfine splittings.

The question is to understand how large mixing angles have to be treated in a model-building point of view. In what follows, we analyze their relationships with hierarchical mass patterns to derive possible hidden components of a unifying structure. We then show how these components can be embedded in a scheme with a replicated gauge group.

2 Schemes with large neutrinos mixings

We start by reconstructing the effective neutrino mass matrix, assuming hierarchical masses, to obtain the following hierarchical structure

$$\begin{pmatrix} \epsilon & \epsilon & \epsilon \\ \epsilon & 1 & 1 \\ \epsilon & 1 & 1 \end{pmatrix} \quad (7)$$

The parameter ϵ is set by the measured ratio of Δm^2 , so $\epsilon^2 \sim 10^{-2}$.

It is worth pointing out that the hierarchy in the neutrino sector is very mild compared to that observed in the charged lepton or in the quark sectors. This observation, together with the appearance of large mixing angles, supports the idea of anarchy as a hypothesis for the description of the neutrino sector. A first task is to compare this possibility to a hierarchical structure.

2.1 Anarchy

Anarchy is basically the idea that there is no fundamental distinction between the three different flavors of neutrinos. The neutrino mass matrix will therefore have no structure

$$\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \quad (8)$$

and the observed values of mixing angles and Δm^2 ratio just reflect Nature's choice of the random $\mathcal{O}(1)$ prefactors that conspired to produce the observed data.

In this scenario, large mixing angles are expected to be generic, so that the small value of θ_{13} puts a severe constraint. Anarchical matrices have been analyzed using statistical tools⁸, and have been shown to be consistent with data with a probability of about 12%.

The anarchical hypothesis can also be embedded in a model⁹ based on an Abelian family symmetry commuting with $SU(5)$, which can reproduce the observed hierarchies among quarks and leptons. This step towards a quark-lepton unification in the Yukawa couplings finds roots in the mass relation $m_b \simeq m_\tau$. However, the agreement in the measured masses between down-type quarks and charged leptons deteriorates for the lighter families. As argued in an earlier work¹⁰, in the framework of Abelian family symmetries, neutrino data could also hint at some quark-lepton unification. In this context, it is natural to ask how large mixing angles can appear in the lepton sector while only small mixing angles are observed in the quark sector. In the case of a family symmetry that commutes with $SU(5)$, the answer is that the charged leptons and the down-type quarks mass matrices are transposed of one another, and therefore, large mixing angles in the lepton sector correspond to hidden, unobservable rotations in the quark sector.

We also note that the anarchical scenario for neutrinos is a possible outcome in a unified scheme with a replicated gauge group¹¹.

2.2 Hierarchy

If the neutrino mass matrix is given by Eq. (7), the values of the mixing angles will also strongly depend on how the prefactor coefficients conspire.

In the generic case, *i.e.* when there is no cancellation between these coefficients, the sub-determinant of the 2-3 block is $D_1 \sim \mathcal{O}(1)$, so we expect the following behavior for eigenmasses and mixing angles,

$$\mu_1 \sim \mathcal{O}(\epsilon), \quad \mu_{2,3} \sim \mathcal{O}(1); \quad \theta_\oplus \sim \mathcal{O}(1), \quad \theta_\ominus, s_{13} \sim \mathcal{O}(\epsilon) \quad (9)$$

If the sub-determinant is suppressed, the solar mixing angle gets enhanced. It becomes large, $\theta_\ominus \sim \mathcal{O}(1)$, for $D_1 \sim \mathcal{O}(\epsilon)$, and close to maximal for $D_1 \sim \mathcal{O}(\epsilon^p)$, with $p > 1$.

The sensitivity of the solar mixing angle to the value of the sub-determinant implies that radiative corrections might play an important role. When the neutrino masses are obtained by the see-saw mechanism, the corrections are indeed important, and depend strongly on the masses of the right-handed neutrinos¹². On the other hand, the mixing angles $\theta_\oplus$ and θ_{13} are more stable.

If radiative corrections are indeed important for the solar mixing angle, low-energy and high-energy parameters will not be simply related. In particular, the correspondence will presumably depend on the unknown masses of the right-handed neutrinos. These difficulties can be avoided in several cases.

For example, in the situation referred to as "right-handed neutrino dominance"¹³, only one right-handed neutrino is responsible for the leading contributions to the low-energy mass matrix

obtained. As an illustration, we take two flavors and

$$m_D = m \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad M^{-1} = \frac{1}{M_0} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad (10)$$

This gives a low-energy neutrino mass matrix with a vanishing determinant

$$m_\nu = m_D M^{-1} m_D^t = \frac{m^2}{M_0} \begin{pmatrix} a^2 & ac \\ ac & c^2 \end{pmatrix} \quad (11)$$

The other right-handed neutrinos give rise to sub-leading contributions. As a result, the sub-determinant can be naturally small, even if radiative corrections are added. Therefore, a natural large solar mixing angle can be obtained if right-handed neutrino dominance holds. However, it is not clear that $\theta_\odot$ will be stable against radiative corrections, because its precise value is a ratio of sub-leading contributions.

Right-handed neutrino dominance is however not a feature of models with $U(1)$ family symmetries of the chiral type. The basic reason is that the right-handed neutrino family charges cancel in the see-saw. The structure of the neutrino matrix is given by the charges of the lepton doublet

$$m_\nu \sim L\bar{N} \cdot \frac{1}{\bar{N}N^t} \cdot (L\bar{N})^t \sim LL^t \quad (12)$$

and the value of the sub-determinant D_1 is not generically suppressed.

2.3 Discrete symmetries

Another way to avoid the sensitivity to radiative corrections is to impose more symmetry. One can think of non-Abelian symmetries, or discrete symmetries as additional components of the flavor structure.

Discrete flavor symmetries in the neutrino mass matrix give the possibility to enforce special values of the mixing angles, as it is suggested by the recent neutrino data. For example, with two generations, a permutation symmetry P_{12} between the two flavors automatically imposes a maximal mixing angle. With three generations, an approximate P_{23} symmetry can give rise to a maximal atmospheric angle, and a large but not maximal solar angle¹⁰.

An extension of a permutation symmetry to three flavors leads to the finite non-Abelian group S_3 of permutations of three objects. It turns out¹⁴ that a tri-bimaximal neutrino mixing can be achieved if neutrinos belong to the (reducible) three dimensional representation of S_3 . The tri-bimaximal mixing corresponds to a bi-maximally mixed neutrino ν_3 and a tri-maximally mixed neutrino ν_2 , corresponding to a maximal atmospheric angle $\theta_\oplus = 45^\circ$, and a large solar angle of about $\theta_\odot \simeq 35^\circ$.

$$U = \begin{pmatrix} \sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{pmatrix} \quad (13)$$

3 Flavor symmetries revisited

Because the Yukawa structure and the mixing angles are very different for quarks and for leptons, it is difficult to embed naturally both sectors in a unified scheme. A hierarchical structure that arises from an Abelian family symmetry can fit nicely the quark sector but leaves the neutrino sector sensitive to order one parameters and to possible radiative corrections. On the other hand, a discrete symmetry that would explain the particular values of the neutrino mixings would require severe fine-tuning to accommodate the hierarchical mass patterns for quarks.

How can Abelian flavor symmetries and discrete flavor symmetries emerge from a common scheme? We contemplate the idea that both components emerge from a replicated gauge structure of the type

$$G_0 = G \times G \times G \quad (14)$$

The fundamental gauge group at the Planck scale contains a copy of the same group G for each generation. Fermions for each family fall into representations of the respective copy of G , but are singlets under the other copies. By symmetry-breaking, the fundamental group is reduced to the diagonal (family-blind) Standard Model gauge group $SU(3) \times SU(2) \times U(1)_Y$. In the chain of symmetry breakings, family-dependent Abelian symmetries are left over at some intermediate step. We assume that they are responsible for the observed hierarchies of the Yukawa couplings. In this framework, the question of family hierarchy reduces to a search for plausible unifying groups and the way to break them. The patterns of the fermion masses and mixings are encoded in the underlying group structure and the breaking path¹¹.

Because the low-energy spectrum contains chiral fermions, a good candidate for a replicated gauge group must be anomaly-free and have complex representations that contain the known fermions. This limits the freedom to the sequence

$$E_6 \supset SO(10) \supset SU(5) \quad (15)$$

It turns out that phenomenologically relevant $U(1)$ symmetries can arise in the breaking

$$E_6 \times E_6 \times E_6 \longrightarrow SU(5) \quad (16)$$

We refer the interested reader to the paper¹¹ for a detailed analysis of the symmetry breaking and the way they can give rise to family-dependent $U(1)$ symmetries. Here, we would only like to point out that this construction leads to Abelian flavor symmetries that are automatically anomaly-free. Moreover, their form is not arbitrary, but is reminiscent of the group structure and the breaking path, although the (finite) number of such possible flavor symmetries turns out to be quite large.

The breaking scheme of Eq. (16) leads to a few viable mass patterns from the point of view of phenomenology. They are expressed by the family charges for the two irreducible representations **10** and $\bar{\mathbf{5}}$ of $SU(5)$ (The family symmetry Y_F commutes with $SU(5)$ because of the breaking scheme that has been chosen). Three models are listed in Table 1. All these models lead to the

	$Y_F(\mathbf{10})$	$Y_F(\bar{\mathbf{5}})$
Model A	(4, 2, 0)	(0, 0, 0)
Model B	(3, 2, 0)	(2, 0, 0)
Model C	(5, 2, 0)	(1, 0, 0)

Table 1: Family charges for three models with phenomenological relevance, obtained in the symmetry breaking $E_6 \times E_6 \times E_6 \rightarrow SU(5)$.

following relations, strongly suggested by data,

$$\frac{m_c}{m_t} \sim \left(\frac{m_s}{m_b}\right)^2 \sim \left(\frac{m_\mu}{m_\tau}\right)^2 \sim \lambda^4 \quad (17)$$

where the expansion parameter λ has a value close to that of the Cabibbo angle $\lambda_c \simeq 0.22$. As a reward, the relations in Eq. (17) automatically give rise to a large mixing angle for the atmospheric neutrinos!

As we can see, these models realize a compromise between the strong hierarchy observed in the quark sector, and the mild hierarchy with large mixing angles in the neutrino sector. In particular, the model A gives rise to an anarchical scenario for the neutrinos. We suggest that a discrete flavor symmetry plays a critical role in this last case. In the replicated gauge structure, it is indeed natural to suppose that the full lagrangian at the Planck scale is invariant under the permutation group S_3 between the three families. In the quark sector, this symmetry is spontaneously broken by the vacuum expectation value of the order parameter, and the $U(1)$ flavor symmetry drives a hierarchical structure. However, the flavor gauge structure being absent for neutrinos, an exact S_3 symmetry can remain. If this is the case, the low-energy neutrino mixing angles that are being measured may be remnants of a replicated structure that holds at a very early stage. A complete understanding of this picture is still in progress¹⁵.

4 Conclusions

The advent of large mixing angles in the neutrino sector brings a new challenge for theorists. While hierarchical structures that arise from $U(1)$ flavor symmetries are well suited to describe the quark sector, their predictions for neutrinos become sensitive to $\mathcal{O}(1)$ coefficients and radiative corrections. Rather, a discrete symmetry based on the group S_3 might be an adequate ingredient chosen by Nature. We have shown that both components arise naturally in a scheme with a replicated gauge group.

Acknowledgments

This work is supported by the United States Department of Energy under grant DE-FG02-97ER41029. The author would like to thank the conference organizers for their hospitality.

References

1. The KamLAND Collaboration, Phys. Rev. Lett. **90** (2003) 021802.
2. The K2K Collaboration, Phys. Rev. Lett. **90** (2003) 041801.
3. M. Apollonio *et al.*, Phys. Lett. B **338** (1998) 383; Phys. Lett. B **420** (1998) 397.
4. The Super-Kamiokande Collaboration, Phys. Rev. Lett. **85** (2000) 3999.
5. The Super-Kamiokande Collaboration, Phys. Lett. B **539** (2002) 179.
6. The SNO Collaboration, Phys. Rev. Lett. **89** (2002) 011301.
7. The LSND collaboration, Phys. Rev. Lett. **77** (1996) 3082; Phys. Rev. Lett. **81** (1998) 1774.
8. G. Altarelli, F. Feruglio and I. Masina, JHEP **0301** (2003) 035; A. de Gouvêa and H. Murayama, hep-ph/0301050.
9. M.S. Berger and K. Siyeon, Phys. Rev. D **63** (2001) 057302.
10. F.-S. Ling and P. Ramond, Phys. Lett. B **543** (2002) 29.
11. F.-S. Ling and P. Ramond, hep-ph/0302264.
12. S. Antusch, J. Kersten, M. Lindner, M. Ratz, Phys. Lett. B **544** (2002) 1; S. Antusch, M. Ratz, JHEP **0211** (2002) 010.
13. S.F. King, JHEP **0209** (2002) 011.
14. P.F. Harrison and W.G. Scott, Phys. Lett. B **557** (2003) 76; J. Kubo, A. Mondragón, M. Mondragón and E. Rodríguez-Jáuregui, hep-ph/0302196.
15. F.-S. Ling, *in preparation*.